

植物生産土壌学

世界の草地土壌写真



No. 23, Haplic Chernozem, Boroll, in Kazakhstan



CHERNOZEMS (C)

Other soils having a mollic A horizon with a moist chroma of 2 or less to a depth of at least 15 cm; having one or more of the following: a calcic or a gypsic horizon, or concentrations of soft powdery lime within 125 cm of the surface when the weighted average textural class is coarse, within 90 cm for medium textures, within 75 cm for fine textures

No.24,
Haplic Chernozem,
Typic Vermudoll,
in Hildesheim, Germany



No. 15,
Gleyic Solonchak,
pink-red of
phenolphthalein
indicates $\text{pH} > 8$



No.17, Orthic Solonetz, Natriboralf
in Tselinograd, Kazakhstan

XEROSOLS (X)

Other soils having a weak ochric A horizon and an aridic moisture regime; lacking permafrost within 200 cm of the surface



No.21,
Calcic Zerosol,
Xerollic Calciorthid
in Konya basin,
central Turkey



No.22, Konya basin, central Turkey,
Vegetation: semi-desert grasses and herbs



Dystric Histosol:

Other Histosols having a pH H_2O , (1:5) of less than 5.5, at least in some part of the soil between 20 and 50 cm from the surface

No. 70,
Dystric Histosol,
Medisaprist,
in Vriezeveen,
The Netherland